Performance

Feeding Directions:

- 1. Before Feeding EquiLix for the first time, animals should be well fed and have access to plenty of fresh water.
- 2. Have one lick block available for every one to five head of horses.
- 3. Provide an adequate source of good quality forage at all times (all they can eat).
- 4. Place lick blocks in stalls or in pastures where horses loaf.
- 5. Make lick blocks available in more than one location in large pastures or where more than one water source is available.
- 6. If animals run out of lick blocks, repeat from Step #1.
- 7. Other salt and minerals may be provided free choice to balance specific rations.
- 8. A two week adaptation period will be required to establish a base consumption rate.
- 9. The above feeding directions are general guidelines and will vary with the overall feeding conditions.

Always use good management practices and adapt to your specific requirements.

Control #:

Guaranteed by: SweetPro Feeds 1505 4th Ave E / Horton, KS 66439 888.229.0475 / www.sweetpro.com

Made in USA

INGREDIENTS: Corn distillers dried grains with solubles, condensed corn distillers solubles, yeast culture, salt, ground flax, pea meal, diatomaceous earth, magnesium oxide, monocalcium phosphate, calcium carbonate, vegetable oil, potassium chloride, garlic, copper proteinate, zinc proteinate, manganese proteinate, copper sulfate, zinc sulfate, manganese sulfate, cobalt carbonate, Ethylenediamine Dihydroiodide, sodium selenite, dried aspergillus oryzae fermentation extract, liquid aspergillus niger fermentation product, vitamin A acetate, vitamin D3 supplement, vitamin E supplement, biotin.

.. 0.50%

. 600 ppm

. 5.0 ppm

. 1200 ppm

.. 750 ppm

.. 8.0 ppm

.. 4.0 ppm

. 0.8 ppm

. 100,000 IU/lb 20,000 IU/lb

......350 IU/lb

.....18 ppm

Guaranteed Analysis

rude Protein, Min	14.00%	Sulfur (S), Min
rude Fat, Min	6.50%	Copper (Cu), Min
rude Fiber, Max	11.00%	Selenium (Se), Min
DF, Max	11.00%	Zinc (Zn), Min
IDF, Max	19.00%	Manganese (Mn), Min
alcium (Ca), Min	1.75%	lodine (I), Min
alcium (Ca), Max	2.25%	Cobalt (Co), Min
hosphorus (P), Min	1.00%	Fluorine, Max
alt (NaCl), Min	7.00%	Vitamin A, Min
alt (NaCl), Max	8.40%	Vitamin D, Min
Magnesium (Mg), Min		Vitamin E, Min
otassium (K), Min	1.10%	Biotin, Min
		0 25 11 4 11/41

Omega 3 Fatty Acid (ALA), Min...... 0.40% Alpha-Amylase (from Aspergillus oryzae).... . Min. 10,180 ug starch hydrolyzed/min/oz. Amyloglucosidase (from Aspergillus niger, Aspergillus oryzae)...... .. Min. 1,540 ug maltose hydrolyzed/min/oz.

CAUTION: Contains 600 ppm copper. Do not feed to sheep!

Net Wt. 50 lbs (22.7 kg)





