Swatho

Reg. No. 620854

Supplement Block for Starting Beef Calves

Net Wt. 102 kg. (225 lbs.)

Guaranteed Analysis

This feed contains added selenium at 5 mg/kg

Crude Protein, Min. 16.00% Crude Fat, Min. 5.00% Crude Fiber, Max. 5.50% Calcium (Ca), Actual 1.25% Phosphorus (P), Actual 1.00% Salt (NaCl), Actual 7.75% Sodium (Na), Actual 3.05% Sulfur (S), Actual 0.75%

Potassium (K), Actual		0.90%
Magnesium (Mg), Actual		
Flourine (F), Max		
Cobalt (Co), Actual	4.0	mg/kg
Copper (Cu), Actual	600	mg/kg
Iodine (I), Actual	8.0	mg/kg
Iron (Fe), Actual	275	mg/kg
Manganese (Mn), Actual	750	mg/kg
Zinc (Zn), Actual	1,200	mg/kg

Vitamin A, Minimum	220,000 IU/kg
Vitamin D, Minimum	44,000 IU/kg
Vitamin E, Minimum	180 IU/kg

Ingredients: Corn Distillers Dried Grains with Solubles, Corn Condensed Distillers Solubles, Salt, Sodium Bicarbonate, Monocalcium Phosphate, Calcium Carbonate, Magnesium Oxide, Potassium Chloride, Yeast Culture Dehydrated, Zinc Sulfate Monohydrate, Manganous Sulfate Monohydrate, Copper Sulfate, Zinc Proteinate (Reg. 990743), Manganese Proteinate (Reg. 990744), Aspergillus Niger Fermentation Product Liquid, Cobalt Carbonate, Ethylenediamine Dihydroiodide, Vitamin A Acetate Vitamin D3 (Reg. 990017), Vitamin E (Reg. 990018).

Starter

Directions for Use

- Before feeding SweetPro® Starter lick blocks for the first time, animals should be well fed.
- 2. Provide access for animals to have plenty of clean, fresh water at all times.
- 3. Have one lick block available per 25 head of cattle.
- 4. Provide an adequate source of good quality roughage and forage at all times (all they can consume).
- 5. Place blocks near water sources or areas cattle frequent regularly.
- 6. Make blocks available in more than one location in large pastures or where more than one water source is available.
- 7. If animals run out of lick blocks, repeat from Step#1.
- Other salt and minerals may be provided free choice to balance specific needs.
- 9. A two week adaptation period will be required to establish a base consumption rate. After adaptation period, recommended intake is 0.34 kgs. per day.

The above feeding directions are general guidelines and will vary with the overall feeding conditions.

Always use good management practices.

Caution!

Contains 600 ppm copper. **DO NOT FEED TO SHEEP!**Do not use in association with another feed containing supplemental selenium.

Control #:

