SUZZE DIO®

Reg. No. 720063

Supplement Block for Sheep

Net Wt. 114 kg. (250 lb.)

Guaranteed Analysis

This feed contains added selenium at 5 mg/kg

Crude Protein, Min	18.00%
Crude Protein from non protein	nitrogen source,
Max	7.00%
Crude Fat, Min	4.00%
Crude Fiber, Max	5.50%
Calcium (Ca), Actual	2.50%
Phosphorus (P), Actual	1.25%
Salt (NaCl), Actual	14.00%
Sodium (Na), Actual	5.60%
Sulfur (S), Actual	1.25%

Potassium (K), Actual	1.10%
Magnesium (Mg), Actual	2.25%
Flourine (F), Max	0.56 mg/kg
Cobalt (Co), Actual	4.0 mg/kg
Copper (Cu), Actual	25 mg/kg
Iodine (I), Actual	8.0 mg/kg
Iron (Fe), Actual	400 mg/kg
Manganese (Mn), Actual	750 mg/kg
Zinc (Zn), Actual	1,200 mg/kg

Vitamin A, Minimum	. 200,000 IU/kg
Vitamin D, Minimum	44,000 IU/kg
Vitamin E, Minimum	180 IU/kg

Ingredients: Com Condensed Distillers Solubles, Com Distillers Dried Grains with Solubles, Salt, Diatomaceous Earth, Magnesium Oxide, Monocalcium Phosphate, Calcium Carbonate, Ammonium SUlfate, Biuret, Calcium Sulfate, Potassium Chloride, Yeast Culture Dehydrated, Zinc Sulfate Monohydrate, Manganous Sulfate Monohydrate, Zinc Proteinate (Reg. 990743), Manganese Proteinate (Reg. 990744), Aspergillus Oryzae Fermentation Extract Dehydrated, Aspergillus Niger Fermentation Product Liquid, Cobalt Carbonate, Sodium Selenite, Ethylenediamine Dihydroiodide, Vitamin Acetate, Vitamin D3 (Reg. # 990017), Vitamin E (Reg. # 990018).



Sheep Block II

Directions for Use

- Before feeding SweetPro Sheep II lick blocks for the first time, animals should be well fed.
- 2. Provide access for animals to have plenty of clean, fresh water at all times.
- 3. Have one lick block available for each 50 head of Sheep.
- 4. Provide an adequate source of good quality roughage and forage at all times (all they can consume).
- 5. Place blocks near water sources or areas Sheep frequent regularly.
- 6. Make blocks available in more than one location in large pastures or where more than one water source is available.
- 7. If animals run out of lick blocks, repeat from Step#1.
- 8. Other salt and minerals may be provided free choice to balance specific needs.
- A two week adaptation period will be required to establish a base consumption rate. After adaptation period, recommended range of intake is 0.11 to 0.14 kgs. per day.

The above feeding directions are general guidelines and will vary with the overall feeding conditions. Always use good management practices.

Caution!

Do not use in association with another feed containing supplemental selenium.

Directions for use must be followed carefully.

Control #:	