

SweetPro[®]

Net Wt. 57 kg. (125 lbs.)



EquiLix
Reg. No. 820772

Guaranteed Analysis

Crude Protein, Minimum.....	13.00%
Crude Fat, Minimum.....	6.00%
Crude Fiber, Maximum.....	11.00%
Calcium (Ca), Minimum.....	2.50%
Phosphorus (P), Minimum.....	1.50%
Salt (NaCl), Minimum.....	12.00%
Sodium (Na), Minimum.....	4.80%
Potassium (K), Minimum.....	1.10%
Magnesium (Mg), Minimum.....	2.50%
Flourine (F), Maximum.....	0.80 mg/kg
Selenium (Se), Minimum.....	5.0 mg/kg
Cobalt (Co), Minimum.....	4.0 mg/kg
Copper (Cu), Minimum.....	600 mg/kg
Iodine (I), Minimum.....	8.0 mg/kg
Iron (Fe), Minimum.....	400 mg/kg
Manganese (Mn), Minimum.....	750 mg/kg
Zinc (Zn), Minimum.....	1,200 mg/kg
Vitamin A, Minimum.....	220,000 IU/kg
Vitamin D, Minimum.....	44,000 IU/kg
Vitamin E, Minimum.....	770 IU/kg

Ingredients: Corn distillers dried grains with solubles, Corn condensed distillers solubles, Yeast Culture Dehydrated, Salt, Linseed Meal, Monocalcium Phosphate, Magnesium Oxide, Calcium Carbonate, Vegetable Oil, Diatomaceous Earth, Calcium Sulfate, Potassium Chloride, Zinc Sulfate Monohydrate, Manganous Sulfate Monohydrate, Copper Sulfate, Zinc Proteinate (Reg. 990743), Manganese Proteinate (Reg. 990744), Aspergillus Oryzae Fermentation Extract Dehydrated, Aspergillus Niger Fermentation Product Liquid, Cobalt Carbonate, Ethylenediamine Dihydroiodide, Vitamin A Acetate, Vitamin D3 (Reg. 990017), Vitamin E (Reg. 990018), d-biotin.

Lot Number #:

A Vitamin and Mineral Supplement for Horses

EquiLix Directions for Use

1. Before Feeding SweetPro EquiLix lick blocks for the first time, animals should be well fed and have access to plenty of clean, fresh water.
2. Have one lick block available for each 25 head of horses.
3. Provide an adequate source of good quality forage at all times (all they can eat).
4. Place lick blocks near areas where animals loaf and at least 30 meters from water.
5. Make lick blocks available in more than one location in large pastures or where more than one water source is available.
6. If animals run out of lick blocks, repeat from Step #1.
7. Provide a fresh, clean supply of water at all times.
8. Other salt and mineral may be provided free choice to balance specific ration requirements.
9. A two week adaptation period will be required to establish a base consumption rate. After adaptation period, consumption will be approximately: 0.25 to 0.5 kgs/day. Monitor horses intake. If exceeds 0.5 kgs check forage quality.

The above feeding directions are general guidelines and will vary with the overall feeding conditions. Always use good management practices.

Caution! Précaution!

Contains 600 mg/kg copper. **DO NOT FEED TO SHEEP!**
 Contient 600 mg/kg de cuivre. **NE PAS NOURRIR LES MOUTONS!**

Do not use in association with another feed containing supplemental selenium.
 Ne pas utiliser en association avec un autre aliment contenant un supplément de sélénium.

Registered by Harvest Fuel, Inc. dba SweetPro Feeds, Walhalla, ND 58282 USA 800-327-9222
 Imported by Nutrition Partners 20 East Avenue, Airdrie, AB T4A 2G8 Canada
 made in U.S.A.