

SweetPro®

Net Wt. 22.7 kg. (50 lbs.)

EquiLix

Guaranteed Analysis

This feed contains added selenium at 5 mg/kg

Crude Protein	Minimum 13.00%
Crude Fat	Minimum 6.00%
Crude Fiber	Maximum 11.00%
Calcium (Ca)	Actual 2.75%
Phosphorus (P)	Actual 1.50%
Salt (NaCl)	Actual 13.00%
Sodium (Na)	Actual 5.20%
Potassium (K)	Actual 1.10%
Magnesium (Mg)	Actual 2.50%
Sulfur (S)	Actual 1.00%
Flourine (F)	Maximum 0.8 mg/kg
Cobalt (Co)	Actual 4.0 mg/kg
Copper (Cu)	Actual 600 mg/kg
Iodine (I)	Actual 8.0 mg/kg
Iron (Fe)	Actual 400 mg/kg
Manganese (Mn)	Actual 750 mg/kg
Zinc (Zn)	Actual 1,200 mg/kg
Vitamin A	Minimum 220,000 IU/kg
Vitamin D	Minimum 44,000 IU/kg
Vitamin E	Minimum 770 IU/kg

Ingredients: Corn distillers dried grains with solubles, Corn condensed distillers solubles, Yeast Culture Dehydrated, Salt, Monocalcium Phosphate, Diatomaceous Earth, Linseed Meal, Pea Meal, Magnesium Oxide, Calcium Carbonate, Canola Oil, Calcium Sulfate Dihydrate, Potassium Chloride, Zinc Sulfate Monohydrate, Manganous Sulfate Monohydrate, Copper Sulfate, Zinc Proteinat (Reg. 990743), Manganese Proteinat (Reg. 990744), Copper Proteinat (Reg. 990741), Aspergillus Oryzae Fermentation Extract Dehydrated, Aspergillus Niger Fermentation Product Liquid, Cobalt Carbonate, Ethylenediamine Dihydroiodide, Vitamin A Acetate, Vitamin D3 (Reg. 990017), Vitamin E (Reg. 990018), d-biotin.

Supplement Block for Horses

EquiLix Directions for Use

1. Before Feeding SweetPro EquiLix lick blocks for the first time, animals should be well fed and have access to plenty of clean, fresh water.
2. Have one lick block available for each 25 head of horses.
3. Provide an adequate source of good quality forage at all times (all they can eat).
4. Place lick blocks near areas where animals loaf and at least 30 meters from water.
5. Make lick blocks available in more than one location in large pastures or where more than one water source is available.
6. If animals run out of lick blocks, repeat from Step #1.
7. Provide a fresh, clean supply of water at all times.
8. Other salt and mineral may be provided free choice to balance specific ration requirements.
9. A two week adaptation period will be required to establish a base consumption rate. After adaptation period, consumption will be approximately: 0.25 to 0.5 kgs/day. Monitor horses intake. If exceeds 0.5 kgs check forage quality.

The above feeding directions are general guidelines and will vary with the overall feeding conditions.
Always use good management practices.

Control #:

Reg. No. 820772



Registered by: Harvest Fuel, Inc., Walhalla, ND 58282 / Horton, KS 66439
800-327-9222 / www.sweetpro.com
made in U.S.A.