

# Mineral EX II



## Mineral EX II Mineral Block for Beef Cattle

Reg. No. 640975

Net Wt: 114 kg (250 lb)

### Guaranteed Analysis

Crude Protein, Min.....	5.25%	Sulfur (S), Actual.....	1.25%
Crude Fat, Min.....	2.00%	Cobalt (Co), Actual.....	25 mg/kg
Crude Fiber, Max.....	2.00%	Copper (Cu), Actual.....	4,200 mg/kg
Calcium (Ca), Actual.....	7.50%	Iodine (I), Actual.....	50 mg/kg
Phosphorus (P), Actual.....	6.20%	Iron (Fe), Actual.....	2,300 mg/kg
Salt (NaCl), Actual.....	14.50%	Manganese (Mn), Actual.....	5,500 mg/kg
Sodium (Na), Actual.....	5.60%	Zinc (Zn), Actual.....	7,200 mg/kg
Magnesium (Mg), Actual.....	3.20%	Selenium (Se), Min.....	30 mg/kg
Vitamin A, Minimum.....	500,000 IU/kg		
Vitamin D, Minimum.....	130,000 IU/kg		
Vitamin E, Minimum.....	500 IU/kg		

**Ingredients:** Monocalcium Phosphate, Corn Distillers Dried Grains with Solubles, Salt, Condensed Corn Distillers Solubles, Calcium Carbonate, Magnesium Oxide, Calcium Sulfate Dihydrate, Potassium Chloride, Yeast Culture Dehydrated, Zinc Sulfate Monohydrate, Manganous Sulfate Monohydrate, Copper Sulfate, Zinc Proteinat (Reg. 990743), Manganese Proteinat (Reg. 990744), Aspergillus Niger Fermentation Product Liquid, Cobalt Carbonate, Sodium Selenite, Ethylenediamine Dihydroiodide, Vitamin A Acetate, Vitamin D3 (Reg. 990017), Vitamin E (Reg. 990018)



Lot Number#: \_\_\_\_\_

### Directions for Use

1. Before feeding SweetPro® Mineral EX II lick blocks for the first time, animals should be well fed.
2. Provide access for animals to have plenty of clean, fresh water at all times.
3. Have one lick block available per 25 head of beef cattle.
4. Provide an adequate source of good quality roughage and forage at all times (all they can consume).
5. Place blocks near water sources or areas cattle frequent regularly.
6. Make blocks available in more than one location in large pastures or where more than one water source is available.
7. If animals run out of lick blocks, repeat from Step#1.
8. Other salt and minerals may be provided free choice to balance specific needs.
9. A two week adaptation period will be required to establish a base consumption rate. After adaptation period, recommended intake is 65 grams per head per day

The above feeding directions are general guidelines and will vary with the overall feeding conditions.

Always use good management practices.

### Caution! Précaution!

Contains 600 mg/kg copper. **DO NOT FEED TO SHEEP!**  
Contient 600 mg/kg de cuivre. **NE PAS DONNER AUX OVINS!**

Do not use in association with another feed containing supplemental selenium.

Ne pas utiliser en association avec un autre aliment contenant du sélénium.

Directions for use must be carefully followed.

Suivre scrupuleusement le mode' emploi.

Registered by Harvest Fuel, Inc. dba SweetPro Feeds, Walhalla, ND 58282 USA (800) 327-9222

Imported by Nutrition Partners 20 East Avenue, Airdrie, AB T4A 2G8 Canada

All rights reserved. SweetPro is a registered trademark of Harvest Fuel, Inc. **Made in U.S.A.** Printed with soy inks.