Swathe

Reg. No. 621334

Supplement Block for Brood Cows & Stocker Cattle on Pasture

(250 lbs.) **Net Wt. 114 kg.**

Guaranteed Analysis

This feed contains added selenium at 5 mg/kg

Crude Protein, Min	16.00%
Crude Fat, Min	5.00%
Crude Fiber, Max	5.50%
Calcium (Ca), Actual	2.00%
Phosphorus (P), Actual	1.25%
Salt (NaCl), Actual	14.25%
Sodium (Na), Actual	5.70%
Sulfur (S), Actual	0.50%

Potassium (K), Actual	1.10%
Magnesium (Mg), Actual	2.25%
Flourine (F), Max	0.56 mg/kg
Cobalt (Co), Actual	4.0 mg/kg
Copper (Cu), Actual	600 mg/kg
Iodine (I), Actual	8.0 mg/kg
Iron (Fe), Actual	400 mg/kg
Manganese (Mn), Actual	750 mg/kg
Zinc (Zn), Actual	1,200 mg/kg

Vitamin A, Minimum	. 200,000 IU/kg
Vitamin D, Minimum	44,000 IU/kg
Vitamin E, Minimum	180 IU/kg

Ingredients: Corn Distillers Dried Grains with Solubles, Condensed Corn Distillers Solubles, Salt, Corn Gluten Meal, Magnesium Oxide, Monocalcium Phosphate, Calcium Carbonate, Canola Oil, Potassium Chloride, Yeast Culture Dehydrated, Zinc Sulfate Monohydrate, Manganous Sulfate Monohydrate, Copper Sulfate, Zinc Proteinate (Reg. 990743), Manganese Proteinate (Reg. 990744), Copper Proteinate (Reg. 990741), Aspergillus Niger Fermentation Product Liquid, Cobalt Carbonate, Sodium Selenite, Ethylenediamine Dihydroiodide, Vitamin A Acetate, Vitamin D3 (Reg. 990017), Vitamin E (Reg. 990018).

SweetPro 16

Directions for Use

- Before feeding SweetPro 16 lick blocks for the first time, animals should be well fed.
- Provide access for animals to have plenty of clean, fresh water at all times.
- 3. Have one lick block available per 25 head of beef cows.
- 4. Provide an adequate source of good quality roughage and forage at all times (all they can consume).
- 5. Place blocks near water sources or areas cattle frequent regularly.
- 6. Make blocks available in more than one location in large pastures or where more than one water source is available.
- 7. If animals run out of lick blocks, repeat from Step#1.
- Other salt and minerals may be provided free choice to balance specific needs.
- 9. A two week adaptation period will be required to establish a base consumption rate. After adaptation period, recommended range of intake is 0.45 kgs. per day.
- Once animals consistently consume more than 0.45 kgs/hd/day, transition to SweetPro FiberMate18. If daily consumption is consistently below 0.45 kgs/hd/day, transition to SweetPro Cattle Kandi.

The above feeding directions are general guidelines and will vary with the overall feeding conditions.

Always use good management practices.

Caution!

Contains 600 mg/kg copper. DO NOT FEED TO SHEEP!

Do not use in association with another feed containing supplemental selenium.

Directions for use must be carefully followed.



Control #:

Registered by Harvest Fuel, Inc. dba SweetPro Feeds, Walhalla, ND 58282 (800) 327-9222