EQUILIX Performance

Feeding Directions:

- Before Feeding EquiLix for the first time, animals should be well fed and have access to plenty of fresh water.
- 2. Have one lick block available for every one to five head of horses.
- **3.** Provide an adequate source of good quality forage at all times (all they can eat).
- 4. Place lick blocks in stalls or in pastures where horses loaf.
- Make lick blocks available in more than one location in large pastures or where more than one water source is available.
- 6. If animals run out of lick blocks, repeat from Step #1.
- **7.** Other salt and minerals may be provided free choice to balance specific rations.
- **8.** A two week adaptation period will be required to establish a base consumption rate.
- **9.** The above feeding directions are general guidelines and will vary with the overall feeding conditions.

Always use good management practices and adapt to your specific requirements.

Control #:

Guaranteed by: SweetPro Feeds 1505 4th Ave E / Horton, KS 66439 888.229.0475 / www.sweetpro.com Made in USA **INGREDIENTS:** Corn distillers dried grains with solubles, condensed corn distillers solubles, yeast culture, salt, ground flax, pea meal, diatomaceous earth, magnesium oxide, monocalcium phosphate, calcium carbonate, vegetable oil, potassium chloride, garlic, copper proteinate, zinc proteinate, manganese proteinate, copper sulfate, zinc sulfate, manganese sulfate, cobalt carbonate, Ethylenediamine Dihydroiodide, sodium selenite, dried aspergillus oryzae fermentation extract, liquid aspergillus niger fermentation product, vitamin A acetate, vitamin D3 supplement, vitamin E supplement, biotin

Guaranteed Analysis

Crude Protein, Min1	4.00%
Crude Fat, Min	6.50%
Omega 3 Fatty Acid (ALA), Min	0.40%
Crude Fiber, Max 1	1.00%
Calcium (Ca), Min	1.75%
Calcium (Ca), Max	2.25%
Phosphorus (P), Min	1.00%
Salt (NaCl), Min	
Salt (NaCl), Max	8.40%
Magnesium (Mg), Min	2.00%
Potassium (K), Min	1.10%
Sulfur (S), Min	0.50%

Copper (Cu), Min	600 ppm
Selenium (Se), Min	5.0 ppm
Zinc (Zn), Min	1200 ppm
Manganese (Mn), Min	750 ppm
lodine (I), Min	8.0 ppm
Cobalt (Co), Min	4.0 ppm
Fluorine, Max	0.8 ppm
Vitamin A, Min	100,000 IU/lb
Vitamin D, Min	20,000 IU/lb
Vitamin E, Min	350 IU/lb
Biotin, Min	18

CAUTION: Contains 600 ppm copper. Do not feed to sheep!

Net Wt. 25 lbs (11.4 kg)









