SWZZLDCO® Net Wt. 250 lbs.

Guaranteed Analysis

Crude Protein	
	(includes not more than 8% equivalent protein from non-protein-nitrogen)
Crude Fat	Minimum 3.50%
Crude Fiber	
Calcium (Ca)	
Phosphorus (P)	
Salt (NaCl)	
Potassium (K)	
Copper (Cu)	Minimum 900 ppm
Zinc (Zn)	
Manganese (Mn)	
Cobalt (Co)	
Iodine (I)	Minimum 12.0 ppm
Vitamin D	
	Aspergillus oryzae)Minimum 3,393 ug starch hydrolyzed/min/oz.
Amyloglucosidase (from Aspergillus niger, Aspergillus oryzae)Minimum 513 ug maltose hydrolyzed/min/oz.	

Ingredients: Corn Distillers Dried Grains with Solubles, Corn Condensed Distillers Solubles, Salt, Calcium Carbonate, Magnesium Oxide, Ammonium Sulfate, Mono-Calcium Phosphate, Grain Product, Biuret, Calcium Sulfate, Calcium Hydroxide, Potassium Chloride, Yeast Culture, Zinc Sulfate, Manganese Sulfate, Copper Sulfate, Zinc Polysaccharide Complex, Zinc Proteinate, Manganese Polysaccharide Complex, Manganese Proteinate, Copper Polysaccharide Complex, Copper Proteinate, Aspergillus Oryzae Extract, Aspergillus Niger Fermentation Liquid, Cobalt Carbonate, Sodium Selenite, Ethylenediamine Dihydroiodide, Vitamin A Acetate (Stability Improved), Vitamin D3 Supplement, Vitamin E Supplement

Zinc, Manganese and Copper are 50% organically complexed.

Does Not Contain Any Animal Protein

Range MaXX

SweetPro® Range MaXX lick blocks are designed for essential fortification of Cow requirements for vitamins A, D & E, plus protein, calcium, phosphorus and trace elements including selenium. Range MaXX should be used for cattle on below average to poor forages. Daily intake will vary depending on forage quality and nutritional needs.

Feeding Directions:

- Before feeding SweetPro® Range MaXX lick blocks for the first time, animals should be well fed.
- 2. Provide access for animals to have plenty of clean, fresh water at all times.
- 3. Provide an adequate source of good quality roughage and forage at all times (all they can consume).
- 4. Have one lick block available per 25 head of beef cattle.
- 5. Place lick blocks near water sources or areas cattle frequent regularly.
- Make lick blocks available in more than one location in large pastures or where more than one water source is available.
- 7. Other salt and minerals may be provided free choice to balance specific needs.
- 8. A two week adaptation period will be required to establish a base consumption rate. After adaptation period, recommended range of intake is .50 to .75 lbs. per head per day.

The above feeding directions are general guidelines and will vary with the overall feeding conditions. Always use good management practices.

Caution!

Contains 900 ppm copper. DO NOT FEED TO SHEEP!

Do not use in association with another feed containing supplemental selenium. This feed shall be used only in accordance with directions furnished on the label.

Control #:_____

