

SweetPro®

(114 kg.) **Net Wt. 250 lbs.**

RangeMaxx
DRY PASTURES / GUT HEALTH

Guaranteed Analysis

Crude Protein	Minimum 40.00%
<small>(includes not more than 8% equivalent protein from non-protein-nitrogen)</small>	
Crude Fat	Minimum 3.50%
Crude Fiber	Maximum 7.50%
Calcium (Ca)	Minimum 4.25% - Maximum 5.00%
Phosphorus (P)	Minimum 1.10%
Salt (NaCl)	Minimum 9.25% - Maximum 10.25%
Magnesium (Mg)	Minimum 4.00%
Potassium (K)	Minimum 0.60%
Sulfur (S)	Minimum 3.50%
Copper (Cu)	Minimum 900 ppm
Selenium (Se)	Minimum 6.75 ppm
Selenium (Se)	Maximum 7.25 ppm
Zinc (Zn)	Minimum 1,400 ppm
Manganese (Mn)	Minimum 1100 ppm
Cobalt (Co)	Minimum 6.0 ppm
Iodine (I)	Minimum 11.00 ppm
Vitamin A	Minimum 150,000 IU/lb.
Vitamin D	Minimum 30,000 IU/lb.
Vitamin E	Minimum 120 IU/lb.
Alpha-Amylase	Minimum 3,393 ug starch hydrolyzed/min/oz.
Amyloglucosidase	Minimum 513 ug maltose hydrolyzed/min/oz.

Ingredients: Corn Distillers Dried Grains with Solubles, Corn Condensed Distillers Solubles, Biuret, Salt, Calcium Carbonate, Magnesium Oxide, Mono-Calcium Phosphate, Yeast Culture, Corn Gluten Meal, Calcium Sulfate, Calcium Hydroxide, Potassium Chloride, Vegetable Oil (Canola and Flax Oil), Zinc Sulfate, Manganese Sulfate, Copper Sulfate, Zinc Polysaccharide Complex, Zinc Proteinate, Manganese Polysaccharide Complex, Manganese Proteinate, Copper Polysaccharide Complex, Copper Proteinate, Cobalt Carbonate, Sodium Selenite, Ethylenediamine Dihydroiodide, Vitamin A Acetate (Stability Improved), Vitamin D3 Supplement, Vitamin E Supplement

Zinc, Manganese and Copper are 50% organically complexed.

Does Not Contain Any Animal Protein

Range Maxx

SweetPro® Range Maxx lick blocks are designed for essential fortification of Cow requirements for vitamins A, D & E, plus protein, calcium, phosphorus and trace elements including selenium. Range Maxx should be used for cattle on below average to poor forages. Daily intake will vary depending on forage quality and nutritional needs.

Feeding Directions:

1. Before feeding SweetPro® Range Maxx lick blocks for the first time, animals should be well fed.
2. Provide access for animals to have plenty of clean, fresh water at all times.
3. Provide an adequate source of good quality roughage and forage at all times (all they can consume).
4. Have one lick block available per 25 head of beef cattle.
5. Place lick blocks near water sources or areas cattle frequent regularly.
6. Make lick blocks available in more than one location in large pastures or where more than one water source is available.
7. Other salt and minerals may be provided free choice to balance specific needs.
8. A two week adaptation period will be required to establish a base consumption rate. After adaptation period, recommended range of intake is .50 to .75 lbs. per head per day.

The above feeding directions are general guidelines and will vary with the overall feeding conditions. Always use good management practices.

CAUTION: USE AS DIRECTED

Contains 900 ppm copper. **DO NOT FEED TO SHEEP!**

Do not use in association with another feed containing supplemental selenium.

Control #: _____



Guaranteed by Harvest Fuel, Inc., Walhalla, ND / Mfd. by SweetPro Feeds, Horton, KS, 66439 / 800.327.9222

All rights reserved. SweetPro is a registered trademark of Harvest Fuel, Inc. Made in U.S.A. Printed with soy inks.

