

SweetPro®

(114 kg.) **Net Wt. 250 lbs.**

Dry Cow Calver-Breeder

Guaranteed Analysis

Crude Protein	Minimum 24.00%
<i>(Includes not more than 10% equivalent protein from non-protein-nitrogen)</i>	
Crude Fat	Minimum 5.00%
Crude Fiber	Maximum 5.50%
ADF	Maximum 9.50%
Calcium (Ca)	Minimum 1.75% - Maximum 2.25%
Phosphorus (P)	Minimum 2.00%
Salt (NaCl)	Minimum 8.00% - Maximum 9.60%
Magnesium (Mg)	Minimum 2.00%
Potassium (K)	Minimum 1.00%
Sulfur (S)	Minimum 1.85%
Copper (Cu)	Minimum 600 ppm
Selenium (Se)	Minimum 4.75 ppm - Maximum 5.25 ppm
Zinc (Zn)	Minimum 1,200 ppm
Manganese (Mn)	Minimum 750 ppm
Iron (Fe)	Minimum 200 ppm
Cobalt (Co)	Minimum 4.0 ppm
Iodine (I)	Minimum 8.0 ppm
Vitamin A	Minimum 100,000 IU/lb.
Vitamin D	Minimum 20,000 IU/lb.
Vitamin E	Minimum 1,100 IU/lb.
Alpha-Amylase (from Aspergillus oryzae).....	Minimum 3,393 ug starch hydrolyzed/min/oz.
Amyloglucosidase (from Aspergillus niger, Aspergillus oryzae).....	Minimum 513 ug maltose hydrolyzed/min/oz.

Ingredients: Corn Distillers Dried Grains with Solubles, Corn Condensed Distillers Solubles, Corn Gluten Meal, Biuret, Ammonium Sulfate, Salt, Mono-Calcium Phosphate, Calcium Sulfate, Magnesium Oxide, Potassium Chloride, Vegetable Oil, Yeast Culture, Zinc Sulfate, Manganese Sulfate, Copper Sulfate, Zinc Polysaccharide Complex, Zinc Proteinate, Manganese Polysaccharide Complex, Manganese Proteinate, Copper Polysaccharide Complex, Copper Proteinate, Dried Aspergillus Oryzae Fermentation Extract, Liquid Aspergillus Niger Fermentation Product, Cobalt Carbonate, Sodium Selenite, Ethylenediamine Dihydroiodide, Vitamin A Acetate (Stability Improved), Vitamin D3 Supplement, Vitamin E Supplement.

Zinc, Manganese and Copper are 50% organically complexed.

Does Not Contain Any Animal Protein

Dry Cow / Calver-Breeder

Dry Cow / Calver-Breeder blocks are designed to meet the nutritional needs of dairy and beef cows prior to calving with emphasis on anionic salts and Vitamin E. For use with dairy cattle upon dry off and segregation. For beef producer best results, use 6 to 8 weeks before calving and 4 to 6 weeks into breeding season.

Feeding Instructions:

1. Before Feeding Dry Cow / Calver-Breeder lick blocks for the first time, animals should be well fed and have access to plenty of clean, fresh water at all times.
2. Have one lick block available for each 25-30 head of beef or dairy cattle.
3. Provide an adequate source of good quality roughage and forage at all times (all they can consume).
4. Move blocks at least 100 feet away from water sources if intake exceeds 1.25 lbs./hd/day.
5. Make blocks available in more than one location in large pastures or where more than one water source is available.
6. If animals run out of lick blocks, repeat from step #1.
7. Other salt and minerals may be provided free choice to balance specific needs.
9. A two week adaptation period will be required to establish a base consumption rate. After adaptation period, recommended range of intake is .50 to 1.25 lbs. per day.

The above feeding directions are general guidelines and will vary with the overall feeding conditions. Always use good management practices.

CAUTION: USE AS DIRECTED

Contains 600 ppm copper. **DO NOT FEED TO SHEEP!**

Do not use in association with another feed containing supplemental selenium.

Control #: _____



Guaranteed by Harvest Fuel, Inc., Walthalla, ND / Mfd. by SweetPro Feeds, Horton, KS, 66439 / 800.327.9222

All rights reserved. SweetPro is a registered trademark of Harvest Fuel, Inc. Made in U.S.A. Printed with soy inks.

