

SweetPro®

(102 kg.) **Net Wt. 225 lbs.**

FIBERMATE 1

REPRODUCTION / RECEIVING / GUT HEALTH

Guaranteed Analysis

Crude Protein	Minimum 20.00%
Crude Fat	Minimum 5.00%
Crude Fiber	Maximum 8.25%
Calcium (Ca)	Minimum 1.25% - Maximum 1.75%
Phosphorus (P)	Minimum 0.60%
Salt (NaCl)	Minimum 8.00% - Maximum 9.00%
Magnesium (Mg)	Minimum 2.20%
Potassium (K)	Minimum 0.50%
Sulfur (S)	Minimum 0.20%
Copper (Cu)	Minimum 600 ppm
Selenium (Se)	Minimum 4.75 ppm - Maximum 5.25 ppm
Zinc (Zn)	Minimum 1,200 ppm
Manganese (Mn)	Minimum 750 ppm
Cobalt (Co)	Minimum 4.0 ppm
Iodine (I)	Minimum 8.0 ppm
Vitamin A	Minimum 100,000 IU/lb.
Vitamin D	Minimum 20,000 IU/lb.
Vitamin E	Minimum 80 IU/lb.

Ingredients: Corn Distillers Dried Grains with Solubles, Corn Condensed Distillers Solubles, Salt, Magnesium Oxide, Yeast Culture, Mono-Calcium Phosphate, Processed Grain By-Products, Calcium Carbonate, Calcium Sulfate, Vegetable Oil (Flax and Canola Oil), Zinc Sulfate, Manganese Sulfate, Copper Sulfate, Zinc Polysaccharide Complex, Zinc Proteinate, Manganese Polysaccharide Complex, Manganese Proteinate, Copper Polysaccharide Complex, Copper Proteinate, Cobalt Carbonate, Ethylenediamine Dihydroiodide, Vitamin A Acetate (Stability Improved), Vitamin D3 Supplement, Vitamin E Supplement.

Zinc, Manganese and Copper are 50% organically complexed.

Does Not Contain Any Animal Protein

FIBERMATE 1

SweetPro® FiberMate 1 lick blocks are designed for essential fortification of Stocker Cattle, Yearling Cattle, and Cow requirements for vitamins A, D & E, plus protein, calcium, phosphorus and trace elements including selenium. FiberMate 1 should be used for cattle on good to excellent forages. Daily intake will vary depending on forage quality and nutritional needs.



Feeding Instructions:

1. Before Feeding SweetPro® FiberMate 1 lick blocks for the first time animals should be well fed.
2. Provide access for animals to have plenty of clean, fresh water at all times.
3. Provide an adequate source of good quality roughage or forage at all times (all they can consume).
4. Have one lick block available per 25 head of beef or dairy cattle.
5. Place lick blocks near water sources or areas cattle frequent regularly.
6. Make lick blocks available in more than one location in large pastures or where more than one water source is available.
7. Other salt and minerals may be provided free choice to balance specific needs.
8. A two week adaptation period will be required to establish a base consumption rate. After adaptation period, recommended range of intake is .75 to 1.25 lbs./hd/day.
9. Once animals consistently consume more than 1.25 lbs./hd/day, transition to FiberMate 2. If daily consumption is consistently below .75 lbs./hd/day, transition to SweetPro® Starter.

The above feeding directions are general guidelines and will vary with the overall feeding conditions. Always use good management practices.

CAUTION: USE AS DIRECTED

Contains 600 ppm copper. **DO NOT FEED TO SHEEP!**

Do not use in association with another feed containing supplemental selenium.

Control #: _____



Guaranteed by Harvest Fuel, Inc., Walhalla, ND / Mfd. by SweetPro Feeds, Horton, KS, 66439 / 800.327.9222

All rights reserved. SweetPro is a registered trademark of Harvest Fuel, Inc. Made in U.S.A. Printed with soy inks.



8 91469 00020 1